



Cold Spring Harbor Symposia on Quantitative Biology

THE STRUCTURE OF DNA

J. D. Watson and F. H. C. Crick

Cold Spring Harb Symp Quant Biol 1953 18: 123-131

Access the most recent version at doi:[10.1101/SQB.1953.018.01.020](https://doi.org/10.1101/SQB.1953.018.01.020)

References

This article cites 21 articles, 3 of which can be accessed free at:
<http://symposium.cshlp.org/content/18/123.refs.html>

Article cited in:

<http://symposium.cshlp.org/content/18/123#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in
the box at the top right corner of the article or [click here](#)

To subscribe to *Cold Spring Harbor Symposia on Quantitative Biology* go to:
<http://symposium.cshlp.org/subscriptions>

THE STRUCTURE OF DNA

J. D. WATSON¹ AND F. H. C. CRICK
 Cavendish Laboratory, Cambridge, England
 (Contribution to the Discussion of Provirus.)

It would be superfluous at a Symposium on Viruses to introduce a paper on the structure of DNA with a discussion on its importance to the problem of virus reproduction. Instead we shall not only assume that DNA is important, but in addition that it is the carrier of the genetic specificity of the virus (for argument, see Hershey, this volume) and thus must possess in some sense the capacity for exact self-duplication. In this paper we shall describe a structure for DNA which suggests a mechanism for its self-duplication and allows us to propose, for the first time, a detailed hypothesis on the atomic level for the self-reproduction of genetic material.

We first discuss the chemical and physical-chemical data which show that DNA is a long fibrous molecule. Next we explain why crystallographic evidence suggests that the structural unit of DNA consists not of one but of two polynucleotide chains. We then discuss a stereochemical model which we believe satisfactorily accounts for both the chemical and crystallographic data. In conclusion we suggest some obvious genetical implications of the proposed structure. A preliminary account of some of these data has already appeared in *Nature* (Watson and Crick, 1953a, 1953b).

I. EVIDENCE FOR THE FIBROUS NATURE OF DNA

The basic chemical formula of DNA is now well established. As shown in Figure 1 it consists of a very long chain, the backbone of which is made up of alternate sugar and phosphate groups, joined together in regular 3' 5' phosphate di-ester linkages. To each sugar is attached a nitrogenous base, only four different kinds of which are commonly found in DNA. Two of these—adenine and guanine—are purines, and the other two—thymine and cytosine—are pyrimidines. A fifth base, 5-methyl cytosine, occurs in smaller amounts in certain organisms, and a sixth, 5-hydroxy-methyl-cytosine, is found instead of cytosine in the T even phages (Wyatt and Cohen, 1952).

It should be noted that the chain is unbranched, a consequence of the regular internucleotide linkage. On the other hand the sequence of the different nucleotides is, as far as can be ascertained, completely irregular. Thus, DNA has some features which are regular, and some which are irregular.

A similar conception of the DNA molecule as a

¹ Aided by a Fellowship from the National Foundation for Infantile Paralysis.

long thin fiber is obtained from physico-chemical analysis involving sedimentation, diffusion, light scattering, and viscosity measurements. These techniques indicate that DNA is a very asymmetrical structure approximately 20 Å wide and many thousands of angstroms long. Estimates of its molecular weight currently center between 5×10^6 and 10^7 (approximately 3×10^4 nucleotides). Surprisingly each of these measurements tend to suggest that the DNA is relatively rigid, a puzzling finding in view of the large number of single bonds (5 per nucleotide) in the phosphate-sugar back-




FIGURE 1. Chemical formula (diagrammatic) of a single chain of desoxyribonucleic acid.

bone. Recently these indirect inferences have been confirmed by electron microscopy. Employing high resolution techniques both Williams (1952) and Kahler *et al.* (1953) have observed, in preparations of DNA, very long thin fibers with a uniform width of approximately 15-20 Å.

II. EVIDENCE FOR THE EXISTENCE OF TWO CHEMICAL CHAINS IN THE FIBER

This evidence comes mainly from X-ray studies. The material used is the sodium salt of DNA (usually from calf thymus) which has been extracted, purified, and drawn into fibers. These fibers are highly birefringent, show marked ultraviolet and infrared dichroism (Wilkins *et al.*, 1951; Fraser and Fraser, 1951), and give good X-ray fiber diagrams. From a preliminary study of these, Wilkins, Franklin and their co-workers at King's College, London (Wilkins *et al.*, 1953; Franklin and Gosling 1953a, b and c) have been able to draw certain general conclusions about the structure of DNA. Two important facts emerge from their work. They are:

(1) *Two distinct forms of DNA exist.* Firstly a crystalline form, Structure A, (Figure 2) which occurs at about 75 per cent relative humidity and contains approximately 30 per cent water. At higher humidities the fibers take up more water, increase in length by about 30 per cent and assume Structure B (Figure 3). This is a less ordered form than Structure A, and appears to be paracrystalline; that is, the individual molecules are all packed parallel to one another, but are not otherwise regularly arranged in space. In Table 1, we have tabulated some of the characteristic features which distinguish the two forms. The transition from A to B is reversible and therefore the two structures are likely to be related in a simple manner.

(2) *The crystallographic unit contains two polynucleotide chains.* The argument is crystallographic and so will only be given in outline. Structure B has a very strong 3.4 Å reflexion on the meridian. As first pointed out by Astbury (1947), this can only mean that the nucleotides in it occur in groups spaced 3.4 Å apart in the fiber direction. On going from Structure B to Structure A the fiber shortens by about 30 per cent. Thus in Structure A the groups must be about 2.5 per cent A apart axially. The measured density of Structure A, (Franklin




FIGURE 2. X-ray fiber diagram of Structure A of deoxyribonucleic acid. (H. M. F. Wilkins and H. R. Wilson, unpub.)

and Gosling, 1953c) together with the cell dimensions, shows that there must be *two* nucleotides in each such group. Thus it is very probable that the crystallographic unit consists of two distinct polynucleotide chains. Final proof of this can only come from a complete solution of the structure.

Structure A has a pseudo-hexagonal lattice, in which the lattice points are 22 Å apart. This distance roughly corresponds with the diameter of fibers seen in the electron microscope, bearing in mind that the latter are quite dry. Thus it is probable that the crystallographic unit and the fiber are the one and the same.

III. DESCRIPTION OF THE PROPOSED STRUCTURE

Two conclusions might profitably be drawn from the above data. Firstly, the structure of DNA is

TABLE 1.
(From Franklin and Gosling, 1953a, b and c)

	Degree of orientation	Repeat distance along fiber axis	Location of first equatorial spacing	Water content	Number of nucleotides within unit cell
Structure A	Crystalline	28 Å	18 Å	30%	22-24
Structure B	Paracrystalline	34 Å	22-24 Å	> 30%	20 (?)




FIGURE 3. X-ray fiber diagram of Structure B of deoxyribonucleic acid. (R. E. Franklin and R. Gosling, 1953a.)

regular enough to form a three dimensional crystal. This is in spite of the fact that its component chains may have an irregular sequence of purine and pyrimidine nucleotides. Secondly, as the structure contains two chains, these chains must be regularly arranged in relation to each other.

To account for these findings, we have proposed (Watson and Crick, 1953a) a structure in which the two chains are coiled round a common axis and joined together by hydrogen bonds between the nucleotide bases (see Figure 4). Both chains follow right handed helices, but the sequences of the atoms in the phosphate-sugar backbones run in opposite directions and so are related by a dyad perpendicular to the helix axis. The phosphates and sugar groups are on the outside of the helix whilst the bases are on the inside. The distance of a phosphorus atom from the fiber axis is 10 Å. We have built our model to correspond to Structure B, which the X-ray data show to have a repeat distance of 34 Å in the fiber direction and a very strong reflexion of spacing 3.4 Å on the meridian of the X-ray pattern. To fit these observations our structure has a nucleotide on each chain every 3.4 Å in the fiber direction, and makes one complete turn after 10 such intervals, i.e., after 34 Å. Our

structure is a well-defined one and all bond distances and angles, including van der Waal distances, are stereochemically acceptable.

The essential element of the structure is the manner in which the two chains are held together by hydrogen bonds between the bases. The bases are perpendicular to the fiber axis and joined together in pairs. The pairing arrangement is very specific, and only certain pairs of bases will fit into the structure. The basic reason for this is that we have assumed that the backbone of each polynucleotide chain is in the form of a regular helix. Thus, irrespective of which bases are present, the glucosidic bonds (which join sugar and base) are arranged in a regular manner in space. In particular, any two glucosidic bonds (one from each chain)




FIGURE 4. This figure is diagrammatic. The two ribbons symbolize the two phosphate-sugar chains and the horizontal rods. The paths of bases holding the chain together. The vertical line marks the fiber axis.

which are attached to a bonded pair of bases, must always occur at a fixed distance apart due to the regularity of the two backbones to which they are joined. The result is that one member of a pair of bases must always be a purine, and the other a pyrimidine, in order to bridge between the two chains. If a pair consisted of two purines, for example, there would not be room for it; if of two pyrimidines they would be too far apart to form hydrogen bonds.

In theory a base can exist in a number of tautomeric forms, differing in the exact positions at which its hydrogen atoms are attached. However, under physiological conditions one particular form of each base is much more probable than any of the others. If we make the assumption that the favored forms always occur, then the pairing requirements are even more restrictive. Adenine can only pair with thymine, and guanine only with cytosine (or 5-methyl-cytosine, or 5-hydroxy-methyl-cytosine). This pairing is shown in detail in Figures 5 and 6. If adenine tried to pair with cytosine it could not form hydrogen bonds, since there would be two hydrogens near one of the bonding positions, and none at the other, instead of one in each.

A given pair can be either way round. Adenine, for example, can occur on either chain, but when it does its partner on the other chain must always be thymine. This is possible because the two glycoside bonds of a pair (see Figures 5 and 6) are symmetrically related to each other, and thus occur in the same positions if the pair is turned over.

It should be emphasized that since each base can form hydrogen bonds at a number of points one can pair up *isolated* nucleotides in a large variety of ways. *Specific* pairing of bases can only be obtained by imposing some restriction, and in our




FIGURE 5. Pairing of adenine and thymine. Hydrogen bonds are shown dotted. One carbon atom of each sugar is shown.




FIGURE 6. Pairing of guanine and cytosine. Hydrogen bonds are shown dotted. One carbon atom of each sugar is shown.

case it is in a direct consequence of the postulated regularity of the phosphate-sugar backbone.

It should further be emphasized that whatever pair of bases occurs at one particular point in the DNA structure, no restriction is imposed on the neighboring pairs, and any *sequence* of pairs can occur. This is because all the bases are flat, and since they are stacked roughly one above another like a pile of pennies, it makes no difference which pair is neighbor to which.

Though any sequence of bases can fit into our structure, the necessity for specific pairing demands a definite relationship between the sequences on the two chains. That is, if we knew the actual order of the bases on one chain, we could automatically write down the order on the other. *Our structure therefore consists of two chains, each of which is the complement of the other.*

IV. EVIDENCE IN FAVOR OF THE COMPLEMENTARY MODEL

The experimental evidence available to us now offers strong support to our model though we should emphasize that, as yet, it has not been proved correct. The evidence in its favor is of three types:

(1) The general appearance of the X-ray picture strongly suggests that the basic structure is helical (Wilkins *et al.*, 1953; Franklin and Gosling, 1953a). If we postulate that a helix is present, we immediately are able to deduce from the X-ray pattern of Structure B (Figure 3), that its pitch is 34 Å and its diameter approximately 20 Å. Moreover, the pattern suggests a high concentration of atoms on the circumference of the helix, in accord with our model which places the phosphate sugar backbone on the outside. The photograph also indicates that the two polynucleotide chains are not

spaced equally along the fiber axis, but are probably displaced from each other by about three-eighths of the fiber axis period, an inference again in qualitative agreement with our model.

The interpretation of the X-ray pattern of Structure A (the crystalline form) is less obvious. This form does not give a meridional reflexion at 3.4 Å, but instead (Figure 2) gives a series of reflexions around 25° off the meridian at spacings between 3 Å and 4 Å. This suggests to us that in this form the bases are no longer perpendicular to the fiber axis, but are tilted about 25° from the perpendicular position in a way that allows the fiber to contract 30 per cent and reduces the longitudinal translation of each nucleotide to about 2.5 Å. It should be noted that the X-ray pattern of Structure A is much more detailed than that of Structure B and so if correctly interpreted, can yield more precise information about DNA. Any proposed model for DNA must be capable of forming either Structure A or Structure B and so it remains imperative for our very tentative interpretation of Structure A to be confirmed.

(2) The anomalous titration curves of undegraded DNA with acids and bases strongly suggests that hydrogen bond formation is a characteristic aspect of DNA structure. When a solution of DNA is initially treated with acids or bases, no groups are titratable at first between pH 5 and pH 11.0, but outside these limits a rapid ionization occurs (Gulland and Jordan, 1947; Jordan, 1951). On back titration, however, either with acid from pH 12 or with alkali from pH 2½, a different titration curve is obtained indicating that the titratable groups are more accessible to acids and bases than is the untreated solution. Accompanying the initial release of groups at pH 11.5 and in the range pH 3.5 to pH 4.5 is a marked fall in the viscosity and the disappearance of strong flow birefringence. While this decrease was originally thought to be caused by a reversible depolymerization (Vilbrandt and Tennent, 1943), it has been shown by Gulland, Jordan and Taylor (1947) that this is unlikely as no increase was observed in the amount of secondary phosphoryl groups. Instead these authors suggested that some of the groups of the bases formed hydrogen bonds between different bases. They were unable to decide whether the hydrogen bonds linked bases in the same or in adjacent structural units. The fact that most of the ionizable groups are originally inaccessible to acids and bases is more easily explained if the hydrogen bonds are between bases within the same structural unit. This point would definitely be established if it were shown that the shape of the initial titration curve was the same at very low DNA concentrations, when the interaction between neighboring structural units is small.

(3) The analytical data on the relative proportion of the various bases show that the amount of adenine is close to that of thymine, and the amount of guanine close to the amount of cytosine + 5-

methyl cytosine, although the ratio of adenine to guanine can vary from one source to another (Chargaff, 1951; Wyatt, 1952). In fact as the techniques for estimation of the bases improve, the ratios of adenine to thymine, and guanine to cytosine + 5-methyl cytosine appear to grow very close to unity. This is a most striking result, especially as the sequence of bases on a given chain is likely to be irregular, and suggests a structure involving paired bases. In fact, we believe the analytical data offer the most important evidence so far available in support of our model, since they specifically support the biologically interesting feature, the presence of complementary chains.

We thus believe that the present experimental evidence justifies the working hypothesis that the essential features of our model are correct and allows us to consider its genetic possibilities.

V. GENETICAL IMPLICATIONS OF THE COMPLEMENTARY MODEL

As a preliminary we should state that the DNA fibers from which the X-ray diffraction patterns were obtained are not artifacts arising in the method of preparation. In the first place, Wilkins and his co-workers (see Wilkins *et al.*, 1953) have shown that X-ray patterns similar to those from the isolated fibers can be obtained from certain intact biological materials such as sperm head and bacteriophage particles. Secondly, our postulated model is so extremely specific that we find it impossible to believe that it could be formed during the isolation from living cells.

A genetic material must in some way fulfil two functions. It must duplicate itself, and it must exert a highly specific influence on the cell. Our model for DNA suggests a simple mechanism for the first process, but at the moment we cannot see how it carries out the second one. We believe, however, that its specificity is expressed by the precise sequence of the pairs of bases. The backbone of our model is highly regular, and the sequence is the only feature which can carry the genetical information. It should not be thought that because in our structure the bases are on the "inside," they would be unable to come into contact with other molecules. Owing to the open nature of our structure they are in fact fairly accessible.

A MECHANISM FOR DNA REPLICATION

The complementary nature of our structure suggests how it duplicates itself. It is difficult to imagine how like attracts like, and it has been suggested (see Pauling and Delbrück, 1940; Friedrich-Freksa, 1940; and Muller, 1947) that self duplication may involve the union of each part with an opposite or complementary part. In these discussions it has generally been suggested that protein and nucleic acid are complementary to each other and that self replication involves the alternate

syntheses of these two components. We should like to propose instead that the specificity of DNA self replication is accomplished without recourse to specific protein synthesis and that each of our complementary DNA chains serves as a template or mould for the formation onto itself of a new companion chain.

For this to occur the hydrogen bonds linking the complementary chains must break and the two chains unwind and separate. It seems likely that the single chain (or the relevant part of it) might itself assume the helical form and serve as a mould onto which free nucleotides (strictly polynucleotide precursors) can attach themselves by forming hydrogen bonds. We propose that polymerization of the precursors to form a new chain only occurs if the resulting chain forms the proposed structure. This is plausible because steric reasons would not allow monomers "crystallized" onto the first chain to approach one another in such a way that they could be joined together in a new chain, unless they were those monomers which could fit into our structure. It is not obvious to us whether a special enzyme would be required to carry out the polymerization or whether the existing single helical chain could act effectively as an enzyme.

DIFFICULTIES IN THE REPLICATION SCHEME

While this scheme appears intriguing, it nevertheless raises a number of difficulties, none of which, however, do we regard as insuperable. The first difficulty is that our structure does not differentiate between cytosine and 5-methyl cytosine, and therefore during replication the specificity in sequence involving these bases would not be perpetuated. The amount of 5-methyl cytosine varies considerably from one species to another, though it is usually rather small or absent. The present experimental results (Wyatt, 1952) suggest that each species has a characteristic amount. They also show that the sum of the two cytosines is more nearly equal to the amount of guanine than is the amount of cytosine by itself. It may well be that the difference between the two cytosines is not functionally significant. This interpretation would be considerably strengthened if it proved possible to change the amount of 5-methyl cytosine in the DNA of an organism without altering its genetical make-up.

The occurrence of 5-hydroxy-methyl-cytosine in the T even phages (Wyatt and Cohen, 1952) presents no such difficulty, since it completely replaces cytosine, and its amount in the DNA is close to that of guanine.

The second main objection to our scheme is that it completely ignores the role of the basic protamines and histones, proteins known to be combined with DNA in most living organisms. This was done for two reasons. Firstly, we can formulate a scheme of DNA reproduction involving it alone and so

from the viewpoint of simplicity it seems better to believe (at least at present) that the genetic specificity is never passed through a protein intermediary. Secondly, we know almost nothing about the structural features of protamines and histones. Our only clue is the finding of Astbury (1947) and of Wilkins and Randall (1953) that the X-ray pattern of nucleoprotamine is very similar to that of DNA alone. This suggests that the protein component, or at least some of it, also assumes a helical form and in view of the very open nature of our model, we suspect that protein forms a third helical chain between the pair of polynucleotide chains (see Figure 4). As yet nothing is known about the function of the protein; perhaps it controls the coiling and uncoiling and perhaps it assists in holding the single polynucleotide chains in a helical configuration.

The third difficulty involves the necessity for the two complementary chains to unwind in order to serve as a template for a new chain. This is a very fundamental difficulty when the two chains are interlaced as in our model. The two main ways in which a pair of helices can be coiled together have been called plectonemic coiling and paranemic coiling. These terms have been used by cytologists to describe the coiling of chromosomes (Huskins, 1941; for a review see Manton, 1950). The type of coiling found in our model (see Figure 4) is called plectonemic. Paranemic coiling is found when two separate helices are brought to lie side by side and then pushed together so that their axes roughly coincide. Though one may start with two regular helices the process of pushing them together necessarily distorts them. It is impossible to have paranemic coiling with two regular simple helices going round the same axis. This point can only be clearly grasped by studying models.

There is of course no difficulty in "unwinding" a single chain of DNA coiled into a helix, since a polynucleotide chain has so many single bonds about which rotation is possible. The difficulty occurs when one has a pair of simple helices with a common axis. The difficulty is a topological one and cannot be surmounted by simple manipulation. Apart from breaking the chains there are only two sorts of ways to separate two chains coiled plectonemically. In the first, one takes hold of one end of one chain, and the other end of the other, and simply pulls in the axial direction. The two chains slip over each other, and finish up separate and end to end. It seems to us highly unlikely that this occurs in this case, and we shall not consider it further. In the second way the two chains must be directly untwisted. When this has been done they are separate and side by side. The number of turns necessary to untwist them completely is equal to the number of turns of one of the chains round the common axis. For our structure this comes to one turn every 34 Å, and thus about 150 turns per million molecular weight of DNA, that is per 5000

A of our structure. The problem of uncoiling falls into two parts:

(1) How many turns must be made, and how is tangling avoided?

(2) What are the physical or chemical forces which produce it?

For the moment we shall be mainly discussing the first of these. It is not easy to decide what is the uninterrupted length of functionally active DNA. As a lower limit we may take the molecular weight of the DNA after isolation, say fifty thousand A in length and having about 1000 turns. This is only a lower limit as there is evidence suggesting a breakage of the DNA fiber during the process of extraction. The upper limit might be the total amount of DNA in a virus or in the case of a higher organism, the total amount of DNA in a chromosome. For T2 this upper limit is approximately 800,000 A which corresponds to 20,000 turns, while in the higher organisms this upper limit may sometimes be 1000 fold higher.

The difficulty might be more simple to resolve if successive parts of a chromosome coiled in opposite directions. The most obvious way would be to have both right and left handed DNA helices in sequence but this seems unlikely as we have only been able to build our model in the right handed sense. Another possibility might be that the long strands of right handed DNA are joined together by compensating strands of left handed polypeptide helices. The merits of this proposition are difficult to assess, but the fact that the phage DNA does not seem to be linked to protein makes it rather unattractive.

The untwisting process would be less complicated if replication started at the ends as soon as the chains began to separate. This mechanism would produce a new two-strand structure without requiring at any time a free single-strand stage. In this way the danger of tangling would be considerably decreased as the two-strand structure is much more rigid than a single strand and would resist attempts to coil around its neighbors. Once the replicating process is started the presence, at the growing end of the pair, of double-stranded structures might facilitate the breaking of hydrogen bonds in the original unduplicated section and allow replication to proceed in a zipper-like fashion.

It is also possible that one chain of a pair occasionally breaks under the strain of twisting. The polynucleotide chain remaining intact could then release the accumulated twist by rotation about single bonds and following this, the broken ends, being still in close proximity, might rejoin.

It is clear that, in spite of the tentative suggestions we have just made, the difficulty of untwisting is a formidable one, and it is therefore worthwhile re-examining why we postulate plectonemic coiling, and not paranemic coiling in which the two helical threads are not intertwined, but merely in close apposition to each other. Our answer is that with

paranemic coiling, the specific pairing of bases would not allow the successive residues of each helix to be in equivalent orientation with regard to the helical axis. This is a possibility we strongly oppose as it implies that a large number of stereochemical alternatives for the sugar-phosphate backbone are possible, an inference at variance to our finding, with stereochemical models (Crick and Watson, 1953) that the position of the sugar-phosphate group is rather restrictive and cannot be subject to the large variability necessary for paranemic coiling. Moreover, such a model would not lead to specific pairing of the bases, since this only follows if the glucosidic links are arranged regularly in space. We therefore believe that if a helical structure is present, the relationship between the helices will be plectonemic.

We should ask, however, whether there might not be another complementary structure which maintains the necessary regularity but which is not helical. One such structure can, in fact, be imagined. It would consist of a ribbon-like arrangement in which again the two chains are joined together by specific pairs of bases, located 3.4 A above each other, but in which the sugar-phosphate backbone instead of forming a helix, runs in a straight line at an angle approximately 30° off the line formed by the pair of bases. While this ribbon-like structure would give many of the features of the X-ray diagram of Structure B, we are unable to define precisely how it should pack in a macroscopic fiber, and why in particular it should give a strong equatorial reflexion at 20-24 A. We are thus not enthusiastic about this model though we should emphasize that it has not yet been disproved.

Independent of the details of our model, there are two geometrical problems which *any* model for DNA must face. Both involve the necessity for some form of super folding process and can be illustrated with bacteriophage. Firstly, the total length of the DNA within T2 is about 8×10^5 A. As its DNA is thought (Siegal and Singer, 1953) to have the same very large M.W. as that from other sources, it must bend back and forth many times in order to fit into the phage head of diameter 800 A. Secondly, the DNA must replicate itself without getting tangled. Approximately 500 phage particles can be synthesized within a single bacterium of average dimensions $10^4 \times 10^4 \times 2 \times 10^4$ A. The total length of the newly produced DNA is some 4×10^8 A, all of which we believe was at some interval in contact with its parental template. Whatever the precise mechanism of replication we suspect the most reasonable way to avoid tangling is to have the DNA fold up into a compact bundle as it is formed.

A POSSIBLE MECHANISM FOR NATURAL MUTATION

In our duplication scheme, the specificity of replication is achieved by means of specific pairing between purine and pyrimidine bases; adenine

with thymine, and guanine with one of the cytosines. This specificity results from our assumption that each of the bases possesses one tautomeric form which is very much more stable than any of the other possibilities. The fact that a compound is tautomeric, however, means that the hydrogen atoms can occasionally change their locations. It seems plausible to us that a spontaneous mutation, which as implied earlier we imagine to be a change in the sequence of bases, is due to a base occurring very occasionally in one of the less likely tautomeric forms, at the moment when the complementary chain is being formed. For example, while adenine will normally pair with thymine, if there is a tautomeric shift of one of its hydrogen atoms it can pair with cytosine (Figure 7). The next time pairing occurs, the adenine (having resumed its more usual tautomeric form) will pair with thymine, but the cytosine will pair with guanine, and so a change in the sequence of bases will have occurred. It would be of interest to know the precise difference in free energy between the various tautomeric forms under physiological conditions.

GENERAL CONCLUSION

The proof or disproof of our structure will have to come from further crystallographic analysis, a




FIGURE 7. Pairing arrangements of adenine before (above) and after (below) it has undergone a tautomeric shift.

task we hope will be accomplished soon. It would be surprising to us, however, if the idea of complementary chains turns out to be wrong. This feature was initially postulated by us to account for the crystallographic regularity and it seems to us unlikely that its obvious connection with self replication is a matter of chance. On the other hand the plectonemic coiling is, superficially at least, biologically unattractive and so demands precise crystallographic proof. In any case the evidence for both the model and the suggested replication scheme will be strengthened if it can be shown unambiguously that the genetic specificity is carried by DNA alone, and, on the molecular side, how the structure could exert a specific influence on the cell.

REFERENCES

- ASTBURY, W. T., 1947, X-Ray Studies of nucleic acids in tissues. *Sym. Soc. Exp. Biol.* 1:66-76.
- CHARGAFF, E., 1951, Structure and function of nucleic acids as cell constituents. *Fed. Proc.* 10:654-659.
- CRICK, F. H. C., and WATSON, J. D., 1953, Manuscript in preparation.
- FRANKLIN, R. E., and GOSLING, R., 1953a, Molecular configuration in sodium thymonucleate. *Nature, Lond.* 171:740-741.
- 1953b, Fiber diagrams of sodium thymonucleate. I. The influence of water content. *Acta Cryst., Camb.* (in press).
- 1953c, The structure of sodium thymonucleate fibers. II. The cylindrically symmetrical Patterson Function. *Acta Cryst., Camb.* (in press).
- FRASER, M. S., and FRASER, R. D. B., 1951, Evidence on the structure of deoxyribonucleic acid from measurements with polarized infra-red radiation. *Nature, Lond.* 167:760-761.
- FRIEDRICH-FREKSA, H., 1940, Bei der Chromosomen Konjugation wirksame Krafte und ihre Bedeutung für die identische Verdopplung von Nucleoproteinen. *Naturwissenschaften* 28:376-379.
- GULLAND, J. M., and JORDAN, D. O., 1946, The macromolecular behavior of nucleic acids. *Sym. Soc. Exp. Biol.* 1: 56-65.
- GULLAND, J. M., JORDAN, D. O., and TAYLOR, H. F. W., 1947, Electrometric titration of the acidic and basic groups of the desoxypentose nucleic acid of calf thymus. *J. Chem. Soc.* 1131-1141.
- HUSKINS, C. L., 1941, The coiling of chromonemata. *Cold Spr. Harb. Symp. Quant. Biol.* 9:13-18.
- JORDAN, D. O., 1951, Physicochemical properties of the nucleic acids. *Prog. Biophys.* 2:51-89.
- KAHLER, H., and LLOYD, B. J., 1953, The electron microscopy of sodium desoxyribonucleate. *Biochim. Biophys. Acta* 10:355-359.
- MANTON, I., 1950, The spiral structure of chromosomes. *Biol. Rev.* 25:486-508.
- MULLER, H. J., 1947, The Gene. *Proc. Roy. Soc. Lond. Ser. B.* 134:1-37.
- PAULING, L., and DEDBRÜCK, M., 1940, The nature of the intermolecular forces operative in biological processes. *Science* 92:77-79.
- SIEGAL, A., and SINGER, S. J., 1953, The preparation and properties of desoxypentosenucleic acid. *Biochim. Biophys. Acta* 10:311-319.
- VILBRANDT, C. F., and TENNENT, H. G., 1943, The effect of

STRUCTURE OF DNA

131

- pH changes upon some properties of sodium thymonucleate solutions. *J. Amer. Chem. Soc.* 63:1806-1809.
- WATSON, J. D., and CRICK, F. H. C., 1953a, A structure for desoxyribose nucleic acids. *Nature, Lond.* 171:737-738.
- 1953b, Genetical implications of the structure of desoxyribose nucleic acid. *Nature, Lond.* (in press).
- WILKINS, M. H. F., GOSLING, R. G., and SEEDS, W. E., 1951, Physical studies of nucleic acids—nucleic acid: an extensible molecule. *Nature, Lond.* 167:759-760.
- WILKINS, M. H. F., and RANDALL, J. T., 1953, Crystallinity in sperm-heads: molecular structure of nucleoprotein in vivo. *Biochim. Biophys. Acta* 10:192 (1953).
- WILKINS, M. H. F., STOKES, A. R., and WILSON, H. R., 1953, Molecular structure of desoxypentose nucleic acids. *Nature, Lond.* 171:738-740.
- WILLIAMS, R. C., 1952, Electron microscopy of sodium desoxyribonucleate by use of a new freeze-drying method. *Biochim. Biophys. Acta* 9:237-239.
- WYATT, G. R., 1952, Specificity in the composition of nucleic acids. In "The Chemistry and Physiology of the Nucleus," pp. 201-213, N. Y. Academic Press.
- WYATT, G. R., and COHEN, S. S., 1952, A new pyrimidine base from bacteriophage nucleic acid. *Nature, Lond.* 170:1072.